

Slow ^4He Quenches Produce Fuzzy, Transient Vortices

R. J. Rivers

Blackett Laboratory, Imperial College, London SW7 2BZ
(February 7, 2008)

We examine the Zurek scenario for the production of vortices in quenches of liquid ^4He in the light of recent experiments. Extending our previous results to later times, we argue that short wavelength thermal fluctuations make vortices poorly defined until after the transition has occurred. Further, if and when vortices appear, it is plausible that they will decay faster than anticipated from turbulence experiments, irrespective of quench rates.

PACS Numbers : 11.27.+d, 05.70.Fh, 11.10.Wx, 67.40.Vs

Several years ago Kibble initiated a programme [1] for examining the consequences of phase transitions in the early universe for large scale structure formation. In particular, cosmic strings produced at these transitions were seen to have the potential for creating the observed fluctuations in the microwave background. Using simple causal arguments [2] he obtained bounds on the density of strings produced in such transitions. Unfortunately such consequences can only be inferred indirectly and with ambiguity. On the other hand, the production and measurement of vortices in condensed matter (the counterpart of cosmic strings in QFT) is a viable experiment. In a series of papers Zurek [3] made predictions for the density of vortices produced during quenches of liquid ^4He and ^3He into their superfluid states, using similar causal arguments.

Zurek's original assumption was that, as we approach the (continuous) transition, the correlation length $\xi(t)$ of the field initially matches the equilibrium correlation length $\xi_{eq}(t)$ of the field. However, once $\xi(t)$ is growing at the speed of (second) sound*, at some time $\bar{t}$ before the transition begins (as characterised by the divergence of $\xi_{eq}(t)$) it freezes. Only after the transition has begun does the field unfreeze, at a comparable time interval $\bar{t}$.

With superfluid He in mind we assume a time-dependent Landau-Ginzburg free energy of the form

$$F(t) = \int d^3x \left(\frac{-\hbar^2}{2m} |\nabla\phi|^2 + \alpha(t)|\phi|^2 + \frac{1}{4}\beta|\phi|^4 \right).$$

In $F(t)$, $\phi = (\phi_1 + i\phi_2)/\sqrt{2}$ is the complex order-parameter field, whose magnitude determines the superfluid density. In equilibrium at temperature T , in a mean field approximation, the chemical potential $\alpha(T/T_c)$ takes the form $\alpha(T/T_c) = \alpha_0\epsilon(T/T_c)$, where $\epsilon = (T/T_c - 1)$ measures the critical temperature T_c relative to T . At a quench the change in T/T_c with time leads most simply to ϵ of the form

$$\epsilon(t) = \epsilon_0 - \frac{t}{\tau_Q}\theta(t)$$

for $-\infty < t < \tau_Q(1 + \epsilon_0)$, after which $\epsilon(t) = -1$. ϵ_0 measures the original relative temperature and τ_Q defines the quench time. The quench begins at time $t = 0$ and the transition from the normal to the superfluid phase begins at time $t = t_0 = \epsilon_0\tau_Q$.

With $\xi_0^2 = \hbar^2/2m\alpha_0$ and $\tau_0 = \hbar/\alpha_0$ setting the fundamental distance and time scales, mean-field critical behaviour fixes [3] $\bar{t} = \sqrt{\tau_0\tau_Q}$. The frozen correlation length of the field $\bar{\xi} = \xi_0(\tau_Q/\tau_0)^{1/4} \approx \xi_{eq}(t_0 - \bar{t}) \approx \xi_{eq}(t_0 + \bar{t})$. The ordering of the field on large scales will be marred by topological defects, in this case vortices. If the *initial* vortex separation is ξ_{def} , then their density, n_{def} , is $n_{def} = O(1/\xi_{def}^2)$. Zurek initially made the *second* assumption (matched by Kibble in QFT) that $\xi_{def} = O(\bar{\xi})$ whereby

$$n_{def} = \frac{1}{f^2\xi_0^2} \sqrt{\frac{\tau_0}{\tau_Q}},$$

where $f = O(1)$. Since ξ_0 also measures cold vortex thickness, $\tau_Q \gg \tau_0$ corresponds to a measurably large number of widely separated vortices.

This is a remarkable prediction in that it says that, in the first instance, the dimensionful interactions $\beta|\phi|^4$ introduce no new scale. It has been the motive for several experiments. One [4] at Lancaster on vortex production in ^4He shows consistency with the prediction, whereas a second experiment [5] on ^4He finds no vortices. On the other hand, vortex production in quenches of ^3He into its *B*-phase in two separate experiments at Grenoble [6] and Helsinki [7] show quantitative agreement with n_{def} above. This is even though it is now argued [8] that the Helsinki experiment should *not* show agreement because of the geometry of the heating event.

In an earlier letter [9] we argued that, RG improvement apart, the simple prediction above for n_{def} is unreliable on two counts. Firstly, and crucially, vortex separation cannot be related simply to field or phase correlation length. It is much more sensible to relate vortex density to the density of line zeroes, n_{zero} , of the field, since

*The corresponding argument in QFT replaces the speed of sound by the speed of light.

a classical vortex is a tube of false vacuum, with a line zero at its core. For Gaussian field fluctuations

$$\langle \phi_a(\mathbf{r}, t) \phi_b(\mathbf{0}, t) \rangle = \delta_{ab} G(\mathbf{r}, t),$$

the line-zero density $n_{zero}(t)$ can be shown [10,11] to have the form (where primes denote differentiation with respect to r)

$$n_{zero}(t) = \frac{-1}{2\pi} \frac{G''(0, t)}{G(0, t)}.$$

It depends manifestly on the *short-distance* behaviour of $G(r, t)$, rather than the correlation length ξ that follows from large-distance behaviour. Secondly, if a line zero is to be counted as a vortex, at the very least it should have fractal dimension close to unity on a scale comparable to the classical vortex thickness ξ_0 , even though it may behave like a random walk on larger scales. Otherwise the density would depend on the scale at which we look. This is equally true for cosmic strings [12].

In [9] we analysed the model until the time that the transition was effected. In this letter we shall now argue that, even when the transition is complete, for slow quenches the short wavelength thermal fluctuations are likely to persist for some time at a level that prevents an approximately scale-invariant definition of vortex density. Further, after that time it is plausible that such vortices as do exist decay more rapidly than is assumed in the analysis of [5].

Motivated by Zurek's later numerical [14,15] simulations, we adopt the time-dependent Landau-Ginzburg (TDLG) equation for F ,

$$\frac{1}{\Gamma} \frac{\partial \phi_a}{\partial t} = -\frac{\delta F}{\delta \phi_a} + \eta_a,$$

where η_a is Gaussian thermal noise, satisfying

$$\langle \eta_a(\mathbf{x}, t) \eta_b(\mathbf{y}', t') \rangle = 2\Gamma T(t) \delta_{ab} \delta(\mathbf{x} - \mathbf{y}') \delta(t - t').$$

In space, time and temperature units in which $\xi_0 = \tau_0 = k_B = 1$, the TDLG equation becomes

$$\dot{\phi}_a(\mathbf{x}, t) = -[-\nabla^2 + \epsilon(t) + \bar{\beta}|\phi(\mathbf{x}, t)|^2]\phi_a(\mathbf{x}, t) + \bar{\eta}_a(\mathbf{x}, t),$$

where $\bar{\beta}$ is the rescaled coupling and $\bar{\eta}$ the rescaled noise. We adopt the approximation of preserving Gaussian fluctuations by linearising the self-interaction as

$$\dot{\phi}_a(\mathbf{x}, t) = -[-\nabla^2 + \epsilon_{eff}(t)]\phi_a(\mathbf{x}, t) + \bar{\eta}_a(\mathbf{x}, t),$$

where ϵ_{eff} contains a (self-consistent) term $O(\bar{\beta}|\phi|^2)$. Additive renormalisation is necessary, so that $\epsilon_{eff} \approx \epsilon$, as given earlier, for $t \leq t_0$.

Self-consistent linearisation is the standard approximation in non-equilibrium QFT [16], but is not strictly necessary here, since numerical simulations that identify line zeroes of the field can be made that use the full self-interaction [15]. However, to date none address the questions we are posing here exactly, and until then there is

virtue in analytic approximations provided they are not taken too seriously.

The solution for $G(r, t)$ from the linear equation for ϕ_a is straightforward, most usefully in the Schwinger proper-time representation, as $G(r, t) =$

$$\int_0^\infty d\tau \bar{T}(t - \tau/2) \left(\frac{1}{4\pi\tau} \right)^{3/2} e^{-r^2/4\tau} e^{-\int_0^\tau ds \epsilon_{eff}(t-s/2)},$$

where $\bar{T}$ is the rescaled temperature.

For constant ϵ_{eff} , as happens at early times, we recover the usual Yukawa correlator. However, a saddle-point calculation at time t_0 gives $\xi(t_0) = O(\xi)$, confirming Zurek's causal result for the freezing of the field correlation length.

A direct substitution of $G(r, t)$ into n_{zero} is impossible because of UV singularities at small τ . However, a starting-point for counting vortices in superfluids is to count line zeroes of an appropriately coarse-grained field, in which structure on a scale smaller than ξ_0 , the classical vortex size, is not present. This is, indeed, the basis of Zurek's and other numerical simulations. For the moment, we put in a momentum cutoff $k^{-1} > l = \bar{l}\xi_0 = O(\xi_0)$ by hand, which corresponds to damping the singularity in $G(r, t)$ at $\tau = 0$ as

$$G_l(r, t) = \int_0^\infty d\tau \frac{\bar{T}(t - \tau/2)}{[4\pi(\tau + l^2)]^{3/2}} e^{-r^2/4\tau} e^{-\int_0^\tau ds \epsilon_{eff}(t-s/2)},$$

making $G_l(0, t)$ finite. We stress that, for $t \approx t_0$, the correlation length ξ remains $O(\xi)$, *independent* of l .

Assuming a *single* zero of $\epsilon_{eff}(t)$ at $t = t_0$, at $r = 0$ the exponential in the integrand peaks at $\tau = \bar{\tau} = 2(t - t_0)$. Expanding about $\bar{\tau}$ to quadratic order gives

$$G_l(0, t) \approx \bar{T}_c e^{2 \int_{t_0}^t du |\epsilon_{eff}(u)|} \int_0^\infty \frac{d\tau e^{-(\tau - 2(t - t_0))^2 |\epsilon'(t_0)|/4}}{[4\pi(\tau + l^2)]^{3/2}}.$$

The effect of the back-reaction is to stop the growth of $G_l(0, t) - G_l(0, t_0) = \langle |\phi|^2 \rangle_t - \langle |\phi|^2 \rangle_0$ at its symmetry-broken value $\bar{\beta}^{-1}$ in our dimensionless units. We see that this cessation of growth requires $\lim_{u \rightarrow \infty} \epsilon_{eff}(u) = 0$. That is, we must choose

$$\epsilon_{eff}(t) = \epsilon(t) + \bar{\beta}(G_l(0, t) - G_l(0, 0)),$$

thereby preserving Goldstone's theorem.

Beyond that, what is remarkable in this approximation is that the density of line zeroes uses *no* property of the self-mass contribution to $\epsilon_{eff}(t)$, self-consistent or otherwise. In n_{zero} all prefactors cancel[†], to give

$$n_{zero}(t) = \frac{1}{4\pi} \frac{\int_0^\infty \frac{d\tau}{(\tau + l^2)^{5/2}} e^{-(\tau - 2(t - t_0))^2/4l^2}}{\int_0^\infty \frac{d\tau}{(\tau + l^2)^{3/2}} e^{-(\tau - 2(t - t_0))^2/4l^2}}$$

[†]Our ignoring prefactors in [9] was fortuitous, leaving our conclusions obtained there unaffected.

on using the definition $\tau_Q = \bar{t}^2$ in natural units. At $t = t_0$ both numerator and denominator are dominated by the short wavelength fluctuations at small τ . Even though the field is correlated over a distance $\bar{\xi} \gg l$ the density of line zeroes $n_{zero} = O(l^{-2})$ depends entirely on the scale at which we look. However, as time passes the peak of the exponential grows and n_{zero} becomes increasingly insensitive to l . How much time we have depends on the magnitude of $\bar{\beta}$, since once $G(0, t)$ has reached this value it stops growing. Since $G(0, t) = O(\exp(((t - t_0)/\bar{t})^2))$ at early times the backreaction is implemented extremely rapidly. We can estimate the time t^* at which this happens by substituting $\epsilon(u)$ for $\epsilon_{eff}(u)$ in the expression for $G_l(0, t)$ above.

For $t > t^*$ the equation for $n_{zero}(t)$ is not so simple since the estimate above, based on a single isolated zero of $\epsilon_{eff}(t)$, breaks down because of the approximate vanishing of $\epsilon_{eff}(t)$ for $t > t^*$. A more careful analysis shows that $G_l(0, t)$ can be written as

$$G_l(0, t) \approx \int_0^\infty \frac{d\tau \bar{T}(t - \tau/2)}{[4\pi(\tau + \bar{l}^2)]^{3/2}} \bar{G}(\tau, t),$$

where $\bar{G}(\tau, t)$ has the same peak as before at $\tau = 2(t - t_0)$, in the vicinity of which

$$\bar{G}(\tau, t) = e^{2 \int_{t_0}^t du |\epsilon_{eff}(u)|} e^{-(\tau - 2(t - t_0))^2 / 4\bar{t}^2},$$

but $\bar{G}(\tau, t) \cong 1$ for $\tau < 2(t - t^*)$. Thus, for $\tau_Q \gg \tau_0$, $G_l(0, t)$ can be approximately separated as

$$G_l(0, t) \cong G_l^{UV}(t) + G^{IR}(t),$$

where

$$G_l^{UV}(t) = \bar{T}(t) \int_0^\infty d\tau / [4\pi(\tau + \bar{l}^2)]^{3/2}$$

describes the scale-dependent short wavelength thermal noise, proportional to temperature, and

$$G^{IR}(t) = \frac{\bar{T}_c}{(8\pi(t - t_0))^{3/2}} \int_{-\infty}^\infty d\tau \bar{G}(\tau, t)$$

describes the scale-independent, temperature independent, long wavelength fluctuations. A similar decomposition $G_l(0, t) \cong G_l^{UV}(t) + G^{IR}(t)$ can be performed in an obvious way, with $G_l^{IR}(t)/G^{IR}(t) = O(t^{-1})$.

Firstly, suppose that, for $t \geq t^*$, $G^{IR}(t) \gg G_l^{UV}(t)$ and $G_l^{IR}(t) \gg G_l^{UV}(t)$, as would be the case for a temperature quench $\bar{T}(t) \rightarrow 0$. Then, with little thermal noise, we have widely separated line zeroes, with density $n_{def}(t) \approx -G_l^{IR}(t)/2\pi G^{IR}(t)$. With $\partial n_{zero}/\partial l$ small in comparison to n_{zero}/l at $l = \xi_0$ we identify such essentially non-fractal line-zeroes with vortices, and n_{zero} with n_{def} . Of course, we require non-Gaussianity to create true classical energy profiles. Nonetheless, the Halperin-Mazenko result may be well approximated even when the

fluctuations are no longer Gaussian [17]. This is supported by the observation that, once the line zeroes have straightened on small scales at $t > t^*$, the Gaussian field energy, largely in field gradients, is

$$\bar{F} \approx \langle \int_V d^3x \frac{1}{2} (\nabla \phi_a)^2 \rangle = -VG''(0, t),$$

where V is the spatial volume. This matches the energy

$$\bar{E} \approx V n_{def}(t) (2\pi G(0, t)) = -VG''(0, t)$$

possessed by a network of classical global strings with density n_{zero} , in the same approximation of cutting off their logarithmic tails.

For times $t > t^*$, but not too late,

$$n_{def}(t) \approx \frac{\bar{t}}{8\pi(t - t_0)} \frac{1}{\xi_0^2} \sqrt{\frac{\tau_0}{\tau_Q}},$$

the solution to Vinen's equation [18]

$$\frac{\partial n_{def}}{\partial t} = -\chi_2 \frac{\hbar}{m} n_{def}^2,$$

where $\chi_2 = 4\pi$ is a *universal constant*.

This decay law is assumed in the analysis of the Lancaster experiments, in which the density of vortices is inferred from the intensity of the signal of scattered second sound. RG improvement leaves ${}^3\text{He}$ unchanged, but for ${}^4\text{He}$ it redefines χ_2 to $\chi_2(1 - T/T_c)^{-1/3} > \chi_2$. The problem is that this makes an already large χ_2 even larger. In the attenuation of second sound the signal to noise ratio is approximately $O(1/\chi_2 t)$. The empirical value of χ_2 used in the Lancaster experiments is not taken from quenches, but turbulent flow experiments. It is suggested [5] that $\chi_2 \approx 0.005$, a good three orders of magnitude smaller than our prediction above. Although the TDLG theory is not very reliable for ${}^4\text{He}$, if our estimate is sensible it does imply that vortices produced in a *temperature* quench decay much faster than those produced in turbulence. ${}^3\text{He}$ experiments provide no check.

In reality, the situation for the Lancaster ${}^4\text{He}$ experiments is even more complex, since they are *pressure* quenches for which the temperature T is almost *constant* at $T \approx T_c$. Unlike temperature quenches [15,16], thermal fluctuations here remain at full strength. Even for ${}^3\text{He}$, T/T_c never gets very small, and henceforth we take $T = T_c$ in $G_l(0, t)$ above. The necessary *time-independence* of $G^{IR}(t)$ for $t > t^*$ is achieved by taking $\epsilon_{eff}(u) = O(u^{-1})$. In consequence, as t increases beyond t^* the relative magnitude of the UV and IR contributions to $G_l(0, t)$ remains *approximately constant*. Further, since for $t = t^*$,

$$e^{2 \int_{t_0}^t du |\epsilon_{eff}(u)|} e^{-(\Delta t)^2 / \bar{t}^2} \approx 1,$$

this ratio is the ratio at $t = t^*$.

Nonetheless, as long as the UV fluctuations are insignificant at $t = t^*$ the density of line zeroes will

remain largely independent of scale. This follows if $G\mathcal{M}^{IR}(t^*) \gg G\mathcal{M}^{UV}(t^*)$, since $G\mathcal{M}_l(0, t)$ becomes scale-independent later than $G_l(0, t)$. In [9] we showed that this is true provided

$$(\tau_Q/\tau_0)(1 - T_G/T_c) < C\pi^4,$$

where $C = O(1)$ and T_G is the Ginzburg temperature. With $\tau_Q/\tau_0 = O(10^3)$ and $(1 - T_G/T_c) = O(10^{-12})$ this inequality is well satisfied for a linearised TDGL theory for ${}^3\text{He}$ derived[†] from the full TDGL theory [13], but there is no way that it can be satisfied for ${}^4\text{He}$, when subjected to a slow mechanical quench, as in the Lancaster experiment, for which $\tau_Q/\tau_0 = O(10^{10})$, since the Ginzburg regime is so large that $(1 - T_G/T_c) = O(1)$. Effectively, the ${}^4\text{He}$ quench is *nineteen* orders of magnitude slower than its ${}^3\text{He}$ counterpart. It is satisfying to find the Ginzburg regime, originally [1,19] but erroneously thought to determine the initial defect density, but then discarded [2,3], reappear as an essential ingredient in undermining universality.

Thus, for ${}^3\text{He}$, with negligible UV contributions, we estimate the primordial density of vortices as

$$n_{def}(t^*) \approx \frac{\bar{t}}{8\pi(t^* - t_0)} \frac{1}{\xi_0^2} \sqrt{\frac{\tau_0}{\tau_Q}},$$

in accord with the original prediction of Zurek. Because of the rapid growth of $G(0, t)$, $(t^* - t_0)/\bar{t} = p > 1 = O(1)$. With p behaving as $(\ln \bar{\beta})^{1/2}$ there is very little variation. For ${}^3\text{He}$ quenches $p \approx 5$ (and for ${}^3\text{He}$ quenches $p \approx 3$). We note that the factor[§] of $f^2 = 8\pi p$ gives a value of $f = O(10)$, in agreement with the empirical results of [6] and the numerical results of [14,15]**.

However, if the inequality is badly violated, as happens with ${}^4\text{He}$ for slow pressure (but not temperature) quenches, then the density of zeroes $n_{def} = O(l^{-2})$ after t^* depends exactly on the scale l at which we look and they are not candidates for vortices. Since the whole of the quench takes place within the Ginzburg regime this is not implausible. However, it is possible that, even though the thermal noise never switches off, there is no more than a postponement of vortex production, since our approximations must break down at some stage. The best outcome is to assume that the effect of the thermal fluctuations on fractal behaviour is diminished, only leading to a delay in the time at which vortices finally appear. Even if we suppose that n_{def} above is a starting point for calculating the density at later times, albeit with a different t_0 , thereby preserving Vinen's law, we then have

the earlier problem of the large $\chi_2 = O(f^2)$. In the absence of any mechanism to reduce its value drastically, this would make it impossible to see vortices. As a separate observation, we note that the large value of f^2 in the prefactor of n_{def} is, in itself, almost enough to make it impossible to see vortices in ${}^4\text{He}$ experiments, should they be present. This will be pursued elsewhere.

In summary, this work suggests that for slow pressure quenches in ${}^4\text{He}$, we see no well-defined vortices at early times because of thermal fluctuations, and it is plausible that, if we do see them at later times, there are less than we would have expected because of their rapid decay and their initial low density. The situation is different for ${}^3\text{He}$. That the density of such vortices as appear may agree with the Zurek prediction is essentially a consequence of dimensional analysis, given that the main effects of the self-interaction have a tendency to cancel in the counting of vortices, without introducing new scales.

This work is the result of a network supported by the European Science Foundation. I thank Tom Kibble, Grisha Volovik, Nuno Antunes and Eleftheria Kavoussanaki for helpful conversations and analysis.

-
- [1] T.W.B. Kibble, *J. Phys.* **A9**, 1387 (1976).
 - [2] T.W.B. Kibble, *Physics Reports* **67**, 183 (1980).
 - [3] W.H. Zurek, *Physics Reports* **276**, Number 4 (1996), and references therein.
 - [4] P.C. Hendry *et al.*, *Nature* **368**, 315 (1994).
 - [5] M.E. Dodd *et al.*, *Phys. Rev. Lett.* **81**, 3703 (1998), *J. Low Temp. Physics* **15**, 89 (1999).
 - [6] C. Bauerle *et al.*, *Nature* **382**, 332 (1996).
 - [7] V.M.H. Ruutu *et al.*, *Nature* **382**, 334 (1996).
 - [8] I.S. Aranson, N.B. Kopnin and V.M. Vinokur, preprint cond-mat/9905286
 - [9] G. Karra and R.J. Rivers, *Phys. Rev. Lett.* **81** 3707 (1998).
 - [10] B.I. Halperin, published in *Physics of Defects*, proceedings of Les Houches, Session XXXV, 1980, editors Balian, Kléman and Poirier (North-Holland Press, 1981) p.816.
 - [11] F. Liu and G.F. Mazenko, *Phys. Rev.* **B46**, 5963 (1992).
 - [12] G. Karra and R.J. Rivers, *Phys. Lett.* **B414**, 28 (1997)
 - [13] Yu. M. Bunkov and O.D. Timofeevskaya, *Phys. Rev. Lett.* **80**, 4927 (1998).
 - [14] P. Laguna and W.H. Zurek, *Phys. Rev. Lett.* **78**, 2519 (1997).
 - [15] N.D. Antunes, L.M.A. Bettencourt, W.H. Zurek, *Phys. Rev. Lett.* **82**, 2824 (1999), and references therein.
 - [16] D. Boyanovsky, H.J. de Vega and R. Holman, preprint hep-ph/9903534.
 - [17] D. Ibaceta, E. Calzetta, preprint hep-ph/9810301
 - [18] W.F. Vinen, *Proc. Roy. Soc. London*, **A242**, 493 (1957).
 - [19] E. Copeland *et al.*, *Physica* **A158**, 460 (1989)

[†]Ignoring the position-dependent temperature of [8]

[§]An errant factor of 3 appeared in the result of [9]

**The temperature quench of the latter is somewhat different from that considered here, but should still give the same results in this case